arXiv:1005.2019v1 [math.NT] 12 May 2010

# A Representation of Permutations with Full Cycle

Ayça Çeşmelioğlu
Sabancı University
MDBF, Orhanlı, Tuzla
34956 İstanbul, Turkey
cesmelioglu@sabanciuniv.edu

**Abstract**

For $q > 2$, Carlitz proved that the group of permutation polynomials (PPs) over $\mathbb{F}_q$ is generated by linear polynomials and $x^{q-2}$. Based on this result, this note points out a simple method for representing all PPs with full cycle over the prime field $\mathbb{F}_p$, where $p$ is an odd prime. We use the isomorphism between the symmetric group $S_p$ of $p$ elements and the group of PPs over $\mathbb{F}_p$, and the well-known fact that permutations in $S_p$ have the same cycle structure if and only if they are conjugate.

## 1 Introduction

For pseudorandom number generation, one of the important problems is to generate sequences with long periods. A sequence $(s_n)$, which is recursively defined by the formula $s_{n+1} = \psi(s_n)$ with an initial value $s_0$, will be a purely periodic sequence when $\psi$ is a PP over the finite field $\mathbb{F}_q$ with $q = p^r$ elements where $p$ is a prime and $r \geq 1$. The period length of $(s_n)$ equals the length of the cycle in which the initial value $s_0$ lies in the permutation induced by $\psi$. The maximal possible period $q$ for $(s_n)$ will be achieved, if $\psi$ corresponds to a PP with full cycle. Note that we use the term *PP with full cycle* for a PP which induces a full cycle permutation.

For $q \geq 2$, PPs over $\mathbb{F}_q$ form a group under composition and reduction modulo $x^q - x$ and this group is isomorphic to the symmetric group $S_q$. In [1], Carlitz proved that any transposition $(0a)$, for $a \in \mathbb{F}_q^*, q > 2$, can be represented by the polynomial

$$-a^2(((x-a)^{q-2} + a^{-1})^{q-2} - a)^{q-2},$$

which shows that any PP over $\mathbb{F}_q$ is of the form

$$\mathcal{P}_n(x) = (\cdots((a_0x + a_1)^{q-2} + a_2)^{q-2} + \cdots + a_n)q - 2 + a_{n+1} \tag{1}$$

for some $n \geq 1$ and $a_0, a_1, \cdots, a_{n+1} \in \mathbb{F}_q$ with $a_0 \neq 0$ or a linear polynomial $\mathcal{P}_0(x) = cx + d \in \mathbb{F}_q[x]$. The cycle structure of PPs given as in (1) was studied in [2] and also some conditions for obtaining PPs with full cycle were determined.

In this note, we characterize all PPs over $\mathbb{F}_p$ with full cycle, where $p$ is an odd prime, in terms of the representation as in (1). As far as we know, no complete characterization of full cycle permutations was given in terms of PPs before.

For the following parts, we will denote the cycle decomposition of a permutation $\mathcal{P}$, which can be expressed as a product of disjoint cycles as

$$\mathcal{P} = \tau_1^{(1)}\tau_2^{(1)}\cdots\tau_{n_1}^{(1)}\tau_1^{(2)}\tau_2^{(2)}\cdots\tau_{n_2}^{(2)}\cdots\tau_1^{(s)}\tau_2^{(s)}\cdots\tau_{n_s}^{(s)}$$

where $\tau_j^{(i)}$, $1 \le j \le n_i$, is a cycle of length $\ell_i$, by

$$\mathcal{T}(\mathcal{P}) = [n_1 \times \ell_1, n_2 \times \ell_2, ..., n_s \times \ell_s].$$

## 2 Main Result

The result of this note is based on the cycle decomposition of the permutations induced by linear polynomials and the following well-known proposition, for which we refer to Section 4.3 of [3].

**Proposition 2.1.** *Two elements of $S_q$ are conjugate in $S_q$ if and only if they have the same cycle structure.*

Note that for the linear polynomial $\mathcal{P}_0(x) = cx + d \in \mathbb{F}_q[x]$, the cycle decomposition is given by

$$\mathcal{T}(\mathcal{P}_0) = \begin{cases} \left[\frac{q}{p} \times p\right] & \text{if } c = 1 \text{ and } d \neq 0, \\ \left[\frac{q-1}{k} \times k, 1 \times 1\right] & \text{if } c \neq 1, \end{cases} \tag{2}$$

where $k$ denotes the order of $c$ in $\mathbb{F}_q^*$.

In order to use the previous proposition, we need a representation for the PP which induces the inverse of a given permutation. For the linear polynomial $\mathcal{P}_0(x)$, the inverse is given by $\mathcal{P}_0^{-1}(x) = c^{-1}x - c^{-1}d$.

In the following parts, we assume $q = p^r, r \ge 1$, and $q > 2$.

**Lemma 2.2.** *Let $\mathcal{P}_n(x) \in \mathbb{F}_q[x]$ denote the PP*

$$\mathcal{P}_n(x) = (\cdots((a_0x + a_1)^{q-2} + a_2)^{q-2} + \cdots + a_n)^{q-2} + a_{n+1},$$

*where $a_0 \in \mathbb{F}_q^*, a_1, a_2, \cdots, a_{n+1} \in \mathbb{F}_q, n \ge 1$.*
*Then the coefficients of*

$$\mathcal{P}_n^{-1}(x) = (\cdots((b_0x + b_1)^{q-2} + b_2)^{q-2} + \cdots + b_n)^{q-2} + b_{n+1}$$

*are given by*

$$b_k = \begin{cases} -a_0^{-1}a_{n+2-k} & \textit{if n+2-k is odd,} \\ -a_0 a_{n+2-k} & \textit{if n+2-k is even,} \end{cases}$$

*for* $1 \leq k \leq n+1$ *and*

$$b_0 = \begin{cases} a_0 & \textit{if n is odd,} \\ a_0^{-1} & \textit{if n is even.} \end{cases}$$

**Proof**:

Let

$$\mathcal{P}_n^{-1}(x) = (\cdots((b_0x+b_1)^{q-2}+b_2)^{q-2}+\cdots+b_n)^{q-2}+b_{n+1}.$$

Then

$$(\mathcal{P}_n \circ \mathcal{P}_n^{-1})(x) = (\cdots((a_0\mathcal{P}_n^{-1}(x)+a_1)^{q-2}+a_2)^{q-2}+\cdots+a_n)^{q-2}+a_{n+1}$$

$$= (\cdots((a_0(\cdots((b_0x+b_1)^{q-2}+b_2)^{q-2}+\cdots+b_n)^{q-2}+a_0b_{n+1}+a_1)^{q-2}+a_2)^{q-2}+\cdots+a_n)^{q-2}+a_{n+1}.$$

We can write $(\mathcal{P}_n \circ \mathcal{P}_n^{-1})(x)$ as

$$(\cdots(((\cdots((a_0b_0x+a_0b_1)^{q-2}+a_0^{-1}b_2)^{q-2}+\cdots+a_0^{-1}b_n)^{q-2}+a_0b_{n+1}+a_1)^{q-2}+a_2)^{q-2}+\cdots+a_n)^{q-2}+a_{n+1}$$

if $n$ is even, and as

$$(\cdots(((\cdots((a_0^{-1}b_0x+a_0^{-1}b_1)^{q-2}+a_0b_2)^{q-2}+\cdots+a_0^{-1}b_n)^{q-2}+a_0b_{n+1}+a_1)^{q-2}+a_2)^{q-2}+\cdots+a_n)^{q-2}+a_{n+1}$$

when $n$ is odd.
We only consider the case where $n$ is odd since the case when $n$ is even can be proved similarly. If we set

$$b_{n+1} = -a_0^{-1}a_1,\ b_n = -a_0a_2,\ b_{n-1} = -a_0^{-1}a_3, \cdots, b_2 = -a_0^{-1}a_n,\ b_1 = -a_0a_{n+1},\ b_0 = a_0$$

then it is easy to see that

$$(\mathcal{P}_n \circ \mathcal{P}_n^{-1})(x) = x$$

by inserting the coefficients $b_i, i = 0, 1, \cdots, n+1$, to the previous equation.

□

The following theorem gives a complete description of the PPs over $\mathbb{F}_p$ with full cycle.

**Theorem 2.3.** *Let* $p$ *be an odd prime,* $\widetilde{\mathcal{P}}_0(x)$ *denote the linear polynomial of the form* $x + d \in \mathbb{F}_p[x], d \in \mathbb{F}_p^*$*, and* $\widetilde{\mathcal{P}}_{2n}(x), n \geq 1$*, denote a PP of the form*

$$\widetilde{\mathcal{P}}_{2n}(x) = (\cdots(((\cdots((x+a_1)^{p-2}+a_2)^{p-2}+\cdots+a_n)^{p-2}+a_{n+1})^{p-2}-a_n)^{p-2}-\cdots-a_2)^{p-2}-a_1$$

*where* $a_1, a_2, a_3, \cdots, a_{n+1} \in \mathbb{F}_p, a_{n+1} \neq 0$*. The PP* $\mathcal{P}(x) \in \mathbb{F}_p[x]$ *is a full cycle if and only if it has a representation as* $\mathcal{P}(x) = \widetilde{\mathcal{P}}_{2n}(x)$ *for some* $n \geq 0$*.*

**Proof**:

The linear polynomial $x + d \in \mathbb{F}_p[x]$ with $d \neq 0$ corresponds to a permutation with full cycle and one can write any PP with full cycle as a conjugate $(\mathcal{P}_n \circ (x+d) \circ \mathcal{P}_n^{-1})(x)$ by using some $\mathcal{P}_n(x) \in \mathbb{F}_p[x], n \geq 0$.
Here, we only consider the case when $n \geq 1$ is even, the case where $n$ is odd can also be dealt with similarly and the case $n = 0$ gives rise to linear PPs which are a full cycle.
Let

$$\mathcal{P}_n(x) = (\cdots((b_0 x + b_1)^{p-2} + b_2)^{p-2} + \cdots + b_n)^{p-2} + b_{n+1} \in \mathbb{F}_p[x].$$

If $n \geq 2$ is even, then we have

$$\mathcal{P}_n^{-1}(x) = (\cdots((b_0^{-1}x - b_0^{-1}b_{n+1})^{p-2} - b_0 b_n)^{p-2} - \cdots - b_0 b_2)^{p-2} - b_0^{-1}b_1.$$

Therefore

$$\begin{aligned} &(\mathcal{P}_n \circ (x+d) \circ \mathcal{P}_n^{-1})(x) \\ = &(\cdots((b_0(\cdots((b_0^{-1}x - b_0^{-1}b_{n+1})^{p-2} - b_0 b_n)^{p-2} - \cdots - b_0 b_2)^{p-2} - b_1 + b_0 d + b_1)^{p-2} + b_2)^{p-2} \\ &+ \cdots + b_n)^{p-2} + b_{n+1} \\ = &(\cdots(((\cdots((x - b_{n+1})^{p-2} - b_n)^{p-2} - \cdots - b_2)^{p-2} + b_0 d)^{p-2} + b_2)^{p-2} + \cdots + b_n)^{p-2} + b_{n+1}. \end{aligned}$$

Note that the coefficients of the resulting PP satisfy the conditions of the theorem.

Conversely, for any given PP of the form $\widetilde{\mathcal{P}}_{2n}(x)$, we can find the linear polynomial $x + d$ and the PP $\mathcal{P}_n(x)$ such that $\widetilde{\mathcal{P}}_{2n}(x) = \mathcal{P}_n(x) \circ (x+d) \circ \mathcal{P}_n^{-1}(x)$ by tracing back the previous part of the proof. □

**Remark 2.1.** *The kth iterate of* $\widetilde{\mathcal{P}}_{2n}(x) \in \mathbb{F}_p[x]$ *is easily seen to be*

$$\widetilde{\mathcal{P}}_{2n}^{(k)}(x) = (\cdots(((\cdots((x + a_1)^{p-2} + a_2)^{p-2} + \cdots + a_n)^{p-2} + k a_{n+1})^{p-2} - a_n)^{p-2} - \cdots - a_2)^{p-2} - a_1,$$

*giving the elements of the Sylow p-subgroup generated by* $\widetilde{\mathcal{P}}_{2n}(x)$.

In terms of applications, PPs with full cycle are certainly the most interesting case. However by using Proposition 2.1, and the isomorphism between the group of PPs over $\mathbb{F}_q$ and the symmetric group $S_q$, it becomes easy to represent permutations in $S_q$ having the same cycle structure as one of the linear polynomials. We give the representation of such polynomials without a proof, since the calculations are easy but rather long.

**Remark 2.2.** *Let* $\widetilde{\mathcal{P}}_{2n}(x) \in \mathbb{F}_q[x]$ *be the PP*

$$\widetilde{\mathcal{P}}_{2n}(x) = (\cdots((b_0 x + b_1)^{q-2} + b_2)^{q-2} + \cdots + b_{2n})^{q-2} + b_{2n+1},$$

*with the coefficients*

$$b_i = \begin{cases} c & \textit{for } i = 0, \\ c a_i & \textit{for } 1 \leq i \leq n \textit{ and } i \textit{ is odd}, \\ c^{-1} a_i & \textit{for } 1 \leq i \leq n \textit{ and } i \textit{ is even}, \\ a_{n+1} & \textit{for } i = n+1, \\ -a_{2n+2-i} & \textit{for } n+2 \leq i \leq 2n+1, \end{cases} \tag{3}$$

*for some* $c \in \mathbb{F}_q^*$ *and* $a_1, a_2, \cdots, a_{n+1} \in \mathbb{F}_q$.
*By Proposition 2.1 and Lemma 2.2, any permutation of* $S_q$ *having the same cycle decomposition as the linear polynomial* $\mathcal{P}_0(x) = cx + d$ *can be represented as in (3) for some* $n \geq 0$.
*Furthermore, the kth iterate of* $\widetilde{\mathcal{P}}_{2n}(x)$ *is*

$$\widetilde{\mathcal{P}}_{2n}^{(k)}(x) = (\cdots((\beta_0 x + \beta_1)^{q-2} + \beta_2)^{q-2} + \cdots + \beta_{2n})^{q-2} + \beta_{2n+1}$$

*with*

$$\beta_i = \begin{cases} c^k & \text{for } i = 0, \\ c^k a_i & \text{for } 1 \leq i \leq n \text{ and } i \text{ is odd}, \\ c^{-k} a_i & \text{for } 1 \leq i \leq n \text{ and } i \text{ is even}, \\ (1 + c^{-1} + \cdots + c^{-(k-1)}) a_{n+1} & \text{for } i = n+1, \\ -a_{2n+2-i} & \text{for } n+2 \leq i \leq 2n+1, \end{cases}$$

*If* $c = 1$ *and* $\widetilde{\mathcal{P}}_{2n}(x) \neq x$ *then* $\mathcal{T}(\widetilde{\mathcal{P}}_{2n}) = \left[\frac{q}{p} \times p\right]$, *and if* $c \neq 1$, *then* $\mathcal{T}(\widetilde{\mathcal{P}}_{2n}) = \left[\frac{q-1}{k} \times k, 1 \times 1\right]$, *where* $k = ord(c)$, *by (2).*

## 3 Acknowledgement

The author would like to thank Wilfried Meidl for his careful reading and useful comments.

## References

[1] L. Carlitz, “Permutations in a finite field ”, Proc. Amer. Math. Soc. 4, 538 1953.

[2] A. Çeşmelioğlu, W. Meidl, A. Topuzoğlu, “On the cycle structure of permutation polynomials ”, Finite Fields Appl. 14, Issue 3, 593–614.

[3] D. S. Dummit, R. M. Foote, “Abstract Algebra ”, John Wiley and Sons, Inc., 2004, pp. 1–932.